

# Continuous Peridural Anaesthesia - Element of Comfort Associated with Vaginal Birth

M. Moisa<sup>1</sup>,  
Mădălina  
Corina Moisa<sup>1</sup>,  
T. Enache<sup>1</sup>,  
C. Pop<sup>2</sup>,  
Victoria  
Nicolau<sup>1</sup>,  
M. Mitran<sup>1</sup>,  
B. Marinescu<sup>1</sup>

<sup>1</sup>. Clinical Hospital of Obstetrics  
and Gynecology  
"Prof. Panait Sărbu", Bucharest  
<sup>2</sup>. UMF „Carol Davila”  
Bucharest, "Elias" Emergency  
Hospital, Bucharest

## Abstract

**Objectives:** Were assessed the effects of the epidural analgesia upon the way of birth appreciated as spontaneous/instrumental birth/caesarean section, both globally and comparative between ropivacaine (R) and bupivacaine (B). Were assessed also the effects of the local anaesthetics used (R and B) on foetal outcome. **Material and method:** After the obtaining of the consent of the Hospital Ethic Committee, we studied a group of 430 parturients, 200 in group R and 230 in group B. Patients were primigravidas or multiparas, ASA I-II, age, degree of maternal block the way of birth. **Results:** The quality of analgesia showed no significant differences between group R and B. Motor block was more intense in group B. Spontaneous birth 77%, out of which 51% needed oxytocic perfusion, vidextractor 15.9%, forceps 3.6%, CS 3%. Fetal assessment showed Apgar score 10 in 38% cases, and 9 in 51% cases. The administration of ropivacaine determined a less intense bloc, allowing the more active participation of the parturient, resulting a higher incidence of spontaneous vaginal births. The versatility offered by CPA may fulfil the variable demands which appear in the labor dynamics, in spontaneous vaginal birth, the instrumental one or CS. **Conclusions:** The minimal impact on foetal outcome, the low cardio and neurotoxic level, the minimal motor block recommends Ropivacaine as local anaesthetic, the most suitable for CPA at birth.

**Keywords:** birth, pain, peridural, Ropivacaine

## Introduction

Pain at birth, even though is considered physiological; it determines a stress response with adrenergic hyperactivity:

- increasing the systemic vascular resistance - lowering the uteroplacental flow;
- decreasing Pa CO<sub>2</sub> - decreasing the foetal O<sub>2</sub>;
- decreasing Pa O<sub>2</sub> - hypoxia + foetal acidosis;
- disrhythmic uterine contractions.

The segmentary CPA concept was created a few decades before and consists in achieving a selective analgesia T10-L1, conserving the sensitive perception and abdomino-pelvin and muscular tone.

The manipulation of the anaesthetic doses of local anaesthetic plus opioid with much more pharmacological subtlety permits the conservation of the pushing reflex and the pelvic diaphragm tone with cu minimal impact on obstetrical outcome.

In spite of these anaesthetic subtleties, the association forceps - epidural persists in many people's minds continues to be perpetuated in books and it

is the first on the few objections list which still feeds the controversy between obstetricians and anaesthesiologists.

The objective of this prospective study consists in:

- the evaluation of the CPA effects of the epidural analgesia upon the way of birth appreciated as spontaneous/instrumental birth/caesarean section, both globally and comparative between ropivacaine (R) and bupivacaine (B);
- the effects of the local anaesthetics used (R and B) on foetal outcome evaluated with APGAR score and NACS.

### Lot's demography

We selected 430 patients whereby:

- 200 in the Ropivacaine group;
- 230 in the Bupivacaine group.

We must point out that all these patients were primiparas or secundiparas ASA I-II, aged 18-40 years, with term or preterm (4) pregnancies, in the active phase of labour, spontaneous or pharmacologically induced, cu with singleton or twin pregnancies (3 cases).

### Indications for CPA

1. both for necessity or therapeutic reasons in:
  - soft tissue distocia;
  - associated with labour test in context of preeclampsia;
  - in preterm birth (4 cases);
  - in association with forceps delivery.
2. as an element of comfort for the parturient, on the patient's request.

We insist on the efficiency and comfort obtained by insertion of the waiting catheter, comparing with analgesia installed after the labour has started for a few hours.

The analgesia must also be efficient until the deliverance of the placenta and the episioraphy are done.

The induction or the stimulation of labour with oxi-tocin has been realised according to the obstetrical protocol of the hospital.

### Monitoring:

- maternal and foetal cardiovascular response: BP, pulse, FCB are monitored with no exception;
- Sp O<sub>2</sub> and ECG are not usually performed;
- tachogram is monitored routinely; in most of the cases we may find a decrease in the intensity of the contractions for maximum 15 minutes.

### Evaluation:

- the quality of analgesia is evaluated after NRS (numeric score of evaluation of pain from 0-10);
- motor block is evaluated after modified Bromaje scale (de la 0-3);
- foetal outcome is evaluated after:
  - ✓ APGAR score at 1 and 5 min;
  - ✓ NAC score at 2 hours and 24 hours.

**NACS:** differentiates the CNS depression induced by the anaesthetic drugs to that provoked by traumatic birth or peripartum asphyxia (values of 30-40 ascertain in an unaffected newborn).

### Results:

The quality of analgesia showed no significant differences between group R and B.

- 80% patients - excellent NRS <2;
- 16% patients - good NRS <5;
- 4% patients - week analgesia NRS >7;
- period of time from the injection up to the delivery of the placenta was similar in both groups R and B;
- motor block: scale Bromage 0 for 51% in group R and 42% in B.

### Way of birth:


1. spontaneous birth 77% - 49% without oxicotic uterine stimulation, and 51% with oxicotic uterine stimulation;
2. vacuum extraction 15.9% - 10% in group R and 21% in B;
3. forceps 3.6% only in group B;
4. CS 3% similar in both groups.

### Foetal evaluation:

**APGAR Score:** 10 - 38%, 9 - 51%, 8 - 10%, 7 - 1%

### NACS score:

- NACS media at 2 hours - 36 for both groups but 21.8% R and 25.7% B have NACS <35;
- at 24 hours - NACS significantly higher for group of ropivacaine (R).



| Figure 1

Figure 2 |




Figure 3 |





Figure 4 |



## Conclusion

Versatility offered by CPA can fulfil the variable demanding which appear in dynamics of labour in spontaneous labour, assisted birth or CS, such as:

- Antispastic effect on cervix;
- CPA coordinates the uterine contractions dynamic distocia is present, shortening phase I of labour;
- In association with labour test, CPA differentiates the soft tissue distocia (reducible) and the bone distocia (irreducible);
- Improves the mother's status who conserves the energy and morale for the 2nd stage of birth coordinating efficiently the effort of pushing with every contraction;
- Improves the foetal oxygenation and utero-placental perfusion - useful element in preeclampsia or in placental insufficiency.

The low percentage (19%) of instrumental births in our Clinic, comparing with the one found in the literature (33%), proves once again that the incidence of forceps delivery is associated with obstetrician's policy, who chooses or not to accept a prolonged 2nd stage and not with CPA.

High incidence of CS cannot be attributable to CPA.

### Ropivacaine:

- better analgesia with minimal motor block;
- lesser neuro and cardiotoxic potential;

- minimal impact foetal outcome, it has important qualities which recommend it as the most suitable anaesthetic for CPA at birth. ■

### References

1. Camorcia M., Capogna G., Epidural bupivacaine and ropivacaine in combination with fentanyl in early labour : A randomized trial .European Journal of anaesthesiology 2003; 20:6639.
2. Pedoto V.A., Baroncini S., et al. A prospective randomized double-blind comparison of epidural bupivacaine 0,125% with epidural ropivacaine 0,2% for labor analgesia. Eur. Jour. anaesthesiology 2003; 20: 979-983.
3. Mc. Ciellan K.J., Fauls D., Ropivacaine: an update of its use in regional anaesthesia Drugs 2000; 60:1065-1093.
4. Capogna G., Camorcia M., Columb M.O., Minimum analgesic doses of fentanyl for epidural analgesia in the first stage of labor . Anesthesia Analgesia 2003; 96: 1178-1182.
5. Simpson D., Curran M.P., Oldfield V., Keating G.M., Ropivacaine: a review of its use in regional anaesthesia and acute pain management. Drugs 2005; 65: 2675-2717.
6. Camorcia M., Capogna G., Lyons G., Columb M., Epidural test dose with bupivacaine and ropivacaine:determination of ED(50) Motor block after epidural administration. British Journal of anaesthesiology 2004; 92: 850-853.
7. Dickenson A.H., Epidural cord pharmacology of pain. British Journal of anesthesiology 1995; 75; 193-200.
8. Sabanethon S., Richardson. Management of pain in labour. British Journal hosp. med. 2005; 50.

Reclamă G21(3)0202 ▼

## Inflamații vulvo-vaginale, un singur răspuns!

  
**TANTUM  
ROSA®**  
Clorhidrat de benzidamină

### O singură substanță activă, 4 acțiuni terapeutice locale:

- **Antiinflamator**
- **Anestezic**
- **Analgezic**
- **Antimicrobian\***



Respectă flora vaginală normală

\*Journal of Chemotherapy vol. 4, n 6 (347-352) 1992, Antimicrobial Activity of Benzidamine, a Non-Steroid Anti-inflammatory Agent N.H. FANAKI M.A EL-NAKEEB.

Tantum Rosa conține clorhidrat de benzidamină și se poate elibera fără prescripție medicală.

Catheter-based neuraxial anesthesia (ie, epidural, CSE, and continuous spinal) allows prolonged anesthesia and titration of the onset of the anesthetic. Single-shot spinal or epidural anesthesia is limited to the duration of action of the injected drug. ANATOMY. Epidural anesthesia is performed by introducing a needle between the lumbar, thoracic, or cervical vertebrae and injecting anesthetic medication into the epidural space, via the epidural needle and/or a catheter inserted through the needle into the epidural space (figure 1). Detailed anatomy of the bony spine and vertebrae are discussed. A spinal anaesthesia " injection of anaesthetic directly into the cerebrospinal fluid " is frequently associated with the epidural analgesia for faster effect. In case of caesarean section, performed as an emergency or not, an anaesthesia becomes necessary. If the epidural catheter is in place, it will be used and thus you can fully participate in the birth of your child. However, in case of insufficient anaesthesia or in case of absence of epidural catheter, the anaesthetist might resort to a spinal anaesthesia or to a general anaesthesia, depending on the amount of time available. What incon