

Nutritional Properties of *Abelmoschus Esculentus* as Remedy to Manage Diabetes Mellitus: A Literature Review

Indah Mohd Amin

Universiti Teknologi MARA (UiTM) Malaysia

Abstract. In 2009, Malaysian Diabetes Association reported that diabetes may affect 1.2 million Malaysian and the disease can be developed from as early as seven years old. Many of the side effects of diabetes can be prevented if glucose levels at normal range is being controlled. This includes using natural plants and herbal supplements as the alternative way to manage and control diabetes. In South East Asia region, herbal medicines such as ampalaya leaves are commonly used to treat diabetic patient. There are also other natural plants to manage diabetes such as using *abelmoschus esculentus* (AE). There had been growing interest among researchers to uncover the bioactive and nutritional properties of AE. Thus, this paper provides an extensive literature review to investigate the current study and topics relevant to the nutritional properties of *abelmoschus esculentus* as a remedy to manage diabetes mellitus. This effort is seen towards providing the evidence support to motivate more scientific research to uncover more scientific nutritional properties of *abelmoschus esculentus* that may be benefited by the modern medicine, thus suggesting new potential target for drug discovery.

Keywords: diabetes mellitus, *abelmoschus esculentus*, traditional remedy

1. Introduction

Diabetes can be described as the increase of glucose in the blood. Careful control blood sugars are crucial to prevent diabetes-related complications. Elderly and people with heart disease whom diagnosed as a diabetic can be severely affected. Many of the side effects of diabetes can be prevented if glucose levels at normal range is maintained. According to WHO estimates, over 180 million people worldwide have diabetes and likely to double by 2030 [1]. The most common type of diabetic is type 2 diabetes as it accounts for 85 to 95% of all cases and constitutes the major and growing public health problem [1]. Nowadays, Malaysian people are suffering from diabetes and heart disease as consequence of the bad lifestyle habit. According to Malaysian Diabetes Association, diabetes may affect 1.2 million Malaysian and can be developed as early as seven years old [2]. Physical inactivity and unhealthy dietary habits which resulted in overweight and insulin resistance which are among risk factors for the development of Type 2 diabetes [1]. There are studies that demonstrate the efficacy some of the natural plants and herbal remedies in treating the diabetes. For example, in Phillipines, herbal medicines commonly used to manage blood glucose include; (i) ampalaya leaves, (ii) apdo ng tilapia, (iii) balat ng kamachilli, (iv) caimito fruis, (v) cogon leaves, (vi) eucalyptus leaves, (vii) neem leaves and (viii) ufat ng makahiya [3]. *Abelmoschus esculentus* (AE) is also one the potential natural plant that been used to manage diabetes. Thus, this paper will provides an extensive literature review to investigate current study and relevant topics concerning the nutritional properties of AE as remedy to manage diabetes mellitus. The next section discusses the literature review on various topics on diabetes mellitus, AE and conclusions.

2. Literature Review

This section provides a summary of the literature review on (i) diabetes mellitus, (ii) AE, (iii) scientific study on anti-diabetic properties of medicinal plants from different countries, (iv) nutritional healing properties of AE and (v) biochemical and nutritional composition of AE.

2.1. Diabetes Mellitus (DM)

Diabetes mellitus (DM) is predominantly characterized by abnormal insulin secretion that leads to elevated glucose [4]. In 2007, International Diabetes Federation (IDF) estimates that in the South-East Asia region, 54 million people were diabetic and an additional 63 million adults had Impaired Glucose Tolerance (IGT). At the regional level, the number of people with diabetes is expected to increase by 71% between 2007 and 2025 [1]. There are two main types of diabetes mellitus, Type 1 or insulin-dependent diabetes mellitus (IDDM) and Type II or noninsulin-dependent diabetes mellitus (NIDDM) [4]. There are many clinical and pathophysiologic differences between Type 1 and Type 2 diabetes mellitus (DM) and the contrasting features is clearly seen in Table 1 [5, 6].

Table 1: Contrasting features of Type 1 and Type 2 Diabetes Mellitus [6]

Feature	Type 1 DM	Type 2 DM
Frequency	10-20%	80-90%
Age at onset	Early (below 35 years)	Late (after 40 years)
Type of onset	Abrupt and severe	Gradual and insidious
Weight	Normal	Obese/non-obese
Family history	<20%	About 60%
Genetic locus	Unknown	Chromosome 6
Pathogenesis	Autoimmune destruction of β -Cells	Insulin resistance, impaired insulin secretion
Islet cell antibodies	Yes	No
Blood insulin level	Decreased insulin	Normal or increased insulin
Islet cell changes	Insulinitis, β -cell depletion	No insulinitis, later fibrosis of islets.
Clinical management	Insulin and diet	Diet, exercise, oral drugs, insulin
Acute complications	Ketoacidosis	Hyperosmolar coma

2.2. *Abelmoschus Esculentus* (AE)

Okra (*abelmoschus esculentus*) also known as lady's finger, is flowering plant in the mallow family. Even though, the plant is cultivated in tropical and warm temperate region around the world but the species is still poorly studied [7]. There is not much report available on the bioactive properties of AE despite its wide usage as medicinal plant. The Table 2 shows the scientific studies done on the bioactive properties of AE.

Table 2: Research on bioactive properties of AE

Bioactive properties	Author
Antioxidative effect	[8] [9]
Mucilages effect	[10]
Anticomplementary & hypoglycemic activity	[11]
Hypoglycemic effect	[12]
Antimicrobial effect	[13]
Anticancer	[14]
Antiproliferative and proapoptotic actions	[15]

Table 2 presents the literature on research of bioactive properties of *abelmoschus esculentus*.

2.3. Scientific study on anti-diabetic properties of medicinal plants from different countries

Table 3: Scientific study on anti-diabetic properties of medicinal plants from different countries

Herbs or medicinal plants	Country/Region	Author
<i>Acacia arabica</i> (Lam.), <i>Aegle marmelos</i> (L.), <i>Allium cepa</i> L., <i>Allium sativum</i> L., <i>Aloe vera</i> (L.), <i>Areca catechu</i> L., <i>Artemisia pallens</i> Wall., <i>Annona squamosa</i> L., <i>Andrographis paniculata</i> Nees, <i>Aerva lanata</i> (L.) Juss, <i>Azadirachta indica</i> A. Juss.	India	[16]
<i>Cochlospermum vitifolium</i>	Mexico	[17] [18] [19]
<i>Tournefortia hirsutissima</i> L.		
<i>Acosmium panamense</i>		
<i>Equisetum myriochaetum</i>		
<i>Psacalium decompositum</i> in		
<i>Ibervillea sonorae</i>	Brazil	[20] [21] [22]
<i>Cecropia pachystachya</i>		
<i>Trema micrantha</i>		
<i>Bauhinia cheilandra</i>	Brazil & Argentina	[23]
<i>Smallantus sonchifolius</i>	Brazil & Uruguay	[24]
<i>Witheringia solanacea</i>	Costa Rica	[25]
<i>Lycium barbarum</i>	China	[26] [27]

<i>Eriobotrya japonica</i>		[28]
<i>Taxus yunnanensis</i>		
<i>Polygonatum odoratum</i>		
<i>Cydonia oblonga</i> Mill. , <i>Helianthus tuberosus</i> L., and <i>Allium porrum</i> L.	Turkey	[29] [30]

Table 3 provides a summary of the literature on the anti-diabetic properties investigation from different countries. The following table 4 presents the literature on few of the nutritional healing properties of AE.

2.4. Nutritional Healing Properties of AE

Table 4: Nutritional healing properties of AE

Healing properties	Author
Alkaline reaction, soothes irritated membrane of the intestinal tract, lowering blood sugar, heal burn & any kind of skin rashes	[31]
Mucilaginous texture soak up unhealthy cholesterol, toxin & mucous waste & clean them from the intestinal tract, acts as laxative that can heal ulcer & may reduce acid reflux, promote good cardiovascular & gastrointestinal health, antioxidant & anticancer	[32]

2.5. Biochemical and nutritional composition of AE

Table 5 presents the literature on research of biochemical and nutritional composition of AE.

Table 5: Biochemical and nutritional composition of AE

Biochemical & nutritional composition	Author
Polyphenolic compounds	[33]
Protein & fat	[34]
Protein, oil & gossypol	[35]
Protein, fat & fiber	[36]
Carbohydrate	[37]
Protein, fat, ash, fiber, phosphorous, and iron	[38]
Calcium & iron	[39]

3. Conclusion

This paper provides an extensive literature review to investigate research and topics relevant to the nutritional properties of AE as remedy to manage diabetes mellitus. The effort is seen towards providing evidence support to motivate more scientific research being conducted to uncover more scientific nutritional properties of AE that may be benefited by the modern medicine, thus suggesting new potential target for drug discovery.

4. Acknowledgements

This research is funded by the Excellence Fund of Research Management Institute, Universiti Teknologi MARA (UiTM) Malaysia. The author would like to thank UiTM Malaysia for its financial and support in completing this work.

5. References

- [1] WHO Regional Office for the South-East Asia, *The Health Situation in the South-East Asia Region: 2001-2007*, World Health Organization, 2009.
- [2] *The Report: Malaysia*, 2009, Oxford Business Group.
- [3] T. Dunning, *Complementary therapies and the management of diabetes and vascular disease: a matter of balance*, John Wiley and Sons, 2006.
- [4] B. Flaws and P. Sionneau, *The treatment of modern western medical diseases with Chinese medicine: a textbook and clinical manual*, Blue Poppy Enterprises, Inc, 2002.
- [5] E. Brown, *Pathology: PreTest self-assessment and review*, McGraw-Hill Professional, 2004.
- [6] H. Mohan, *Essential Pathology for Dental Students*, Jaypee Brothers Publishers, 2005

- [7] E. G. Maganha, R. d. C. Halmenschlager, R. M. Rosa, J. A. P. Henriques, A. L. L. d. P. Ramos and J. Saffi, "Pharmacological evidences for the extracts and secondary metabolites from plants of the genus Hibiscus," *Food Chemistry*, vol. 118, no. 1, 2010, pp. 1-10,
- [8] O. E. Adelakun, O. J. Oyelade, B. I. O. Ade-Omowaye, I. A. Adeyemi, M. V. d. Venter, Chemical composition and the antioxidative properties of Nigerian Okra Seed (*Abelmoschus esculentus* Moench) Flour, 2009, 47 (6): 1123-1126.
- [9] F. O. Adetuyi, Antioxidant Degradation in Six Indigenous Okra *Abelmoschus esculentus* (L) Moench Varieties During Storage in Nigeria, *Journal of Food Technology*, 2008, 6 (5): 227-230.
- [10] K. Ameena, C. Dilip, R. Saraswathi, P. N. Krishnan, C. Sankar, S. P. Simi, Isolation of the mucilages from *Hibiscus rosasinensis* linn. and Okra (*Abelmoschus esculentus* linn.) and studies of the binding effects of the mucilages, *Asian Pacific Journal of Tropical Medicine*, 2010, 3(7): 539-543.
- [11] M. Tomoda, N. Shimizu, R. Gonda, M. Kanari, H. Yamada, H. Hiki, Anticomplementary and hypoglycemic activity of Okra and Hibiscus mucilages, *Carbohydrate Research*, 190 (2): 323-328.
- [12] X Zhenzhong, and S. Hongjun, Effects of okra capsule combined with valsartan in treatment of early diabetic nephropathy with microalbuminuria, *Modern Journal of Integrated Traditional Chinese and Western Medicine* 2010, 3.
- [13] C. Lengsfeld, F. Titgemeyer, G. Faller and A. Hensel, Glycosylated Compounds from Okra Inhibit Adhesion of *Helicobacter pylori* to Human Gastric Mucosa, *J. Agric. Food Chem.*, 2004, 52 (6): 1495–1503.
- [14] R. D. Dan and C. Gu, Inhibition Effect of Okra Polysaccharides on Proliferation of Human Cancer Cell Lines, *Food Science* 2010, 21.
- [15] M. Vayssade, N. Sengkhamparn, R. Verhoef, C. Delaigue, O. Oundiam, P. Vigneron, A. G. J. Voragen, H. A. Schols, and M. D. Nagel, Antiproliferative and proapoptotic actions of okra pectin on B16F10 melanoma cells, *Phytotherapy Research*, 2010, 24 (7): 982–989.
- [16] P. K. Mukherjee, K. Maiti, K. Mukherjee and P. J. Houghton. Leads from Indian medicinal plants with hypoglycemic potentials. *Journal of Ethnopharmacology*. 2006, 106 (1): 1-28
- [17] J. C. Sánchez-Salgado, R. R. Ortiz-Andrade, F. Aguirre-Crespo, J. Vergara-Galicia, I. León-Rivera, S. Montes, R. Villalobos-Molina and S. Estrada-Soto, Hypoglycemic, vasorelaxant and hepatoprotective effects of *Cochlospermum vitifolium* (Willd.) Sprengel: A potential agent for the treatment of metabolic syndrome, *Journal of Ethnopharmacology*, 2007, 109 (3): 400-405
- [18] F. J. Alarcon-Aguilar, M. Jimenez-Estrada, R. Reyes-Chilpa and R. Roman-Ramos, Hypoglycemic effect of extracts and fractions from *Psacalium decompositum* in healthy and alloxan-diabetic mice, *Journal of Ethnopharmacology*, 2000, 72 (1-2): 21-27.
- [19] F. J. Alarcon-Aguilar, F. Calzada-Bermejo, E. Hernandez-Galicia, C. Ruiz-Angeles and R. Roman-Ramos, Acute and chronic hypoglycemic effect of *Ibervillea sonorae* root extracts-II, *Journal of Ethnopharmacology*, 2005, 97 (3): 447-452.
- [20] D. M. O. Aragão, L. Guarize, J. Lanini, J. C. D. Costa, R. M. G. Garcia and E. Scio, Hypoglycemic effects of *Cecropia pachystachya* in normal and alloxan-induced diabetic rats, *Journal of Ethnopharmacology*, 2010, 128 (3): 629-633.
- [21] T. Schoenfelder, T. M. Cirimbelli and V.C. Zanette, Acute effect of *Trema micrantha* (Ulmaceae) on serum glucose levels in normal and diabetic rats, *Journal of Ethnopharmacology*, 2006, 107 (3): 456-459.
- [22] E. R. Almeida, M. C. Guedes, J. F. C. Albuquerque and H. Xavier, Hypoglycemic effect of *Bauhinia cheilandra* in rats, *Fitoterapia*, 2006, 77(4): 276-278.
- [23] M. J. Aybar, A. N. S. Riera, A. Grau, and S. S. Sánchez, Hypoglycemic effect of the water extract of *Smallantus sonchifolius* (yacon) leaves in normal and diabetic rats, *Journal of Ethnopharmacology*, 2001, 74 (2): 125-132.
- [24] O. Hnatyszyn, J. Miño, G. Ferraro and C. Acevedo, The hypoglycemic effect of *Phyllanthus sellowianus* fractions in streptozotocin-induced diabetic mice, *Phytomedicine*, 2002, 9 (6): 556-559.
- [25] C. Herrera, P. M. G. Barrantes, F. Binns, M. Vargas, L. Poveda and S. Badilla, Hypoglycemic and antihyperglycemic effect of *Witheringia solanacea* in normal and alloxan-induced hyperglycemic rats, *Journal of*

Ethnopharmacology, 2011, 133 (2): 907-910.

- [26] O. Luo, Y. Cai, J. Yan, M. Sun and H. Corke, Hypoglycemic and hypolipidemic effects and antioxidant activity of fruit extracts from *Lycium barbarum*, *Life Sciences*, 2004, 76 (2): 137-149.
- [27] J. Chen, W. L. Li, J. L. Wu, B. R. Ren and H. Q. Zhang, Hypoglycemic effects of a sesquiterpene glycoside isolated from leaves of loquat (*Eriobotrya japonica* (Thunb.) Lindl.), *Phytomedicine*, 2008, 15 (1-2): 98-102.
- [28] A. H. Banskota, N. T. Nguyen, Y. Tezuka, T. Nobukawa and S. Kadota, Hypoglycemic effects of the wood of *Taxus yunnanensis* on streptozotocin-induced diabetic rats and its active components, *Phytomedicine*, 2006, 13 (1-2): 109-114.
- [29] M. Aslan, N. Orhan, D. D. Orhan and F. Ergun, Hypoglycemic activity and antioxidant potential of some medicinal plants traditionally used in Turkey for diabetes, *Journal of Ethnopharmacology*, 2010, 128 (2): 384-389.
- [30] D. D. Orhan, M. Aslan, N. Sendogdu, F. Ergun and E. Yesilada, Evaluation of the hypoglycemic effect and antioxidant activity of three *Viscum album* subspecies (European mistletoe) in streptozotocin-diabetic rats, *Journal of Ethnopharmacology*, 2005, 98 (1-2): 95-102.
- [31] S. P. Bansal. *Healing Power of Foods*. Pustak Mahal, 2002.
- [32] E. M. Collins. *An A-Z Guide to Healing Foods: A Shopper's Reference*. Conari, 2010.
- [33] P. Arapitsas, Identification and quantification of polyphenolic compounds from okra seeds and skins, *Food Chemistry*, 2008, 110 (4): 1041-1045.
- [34] O. J. Oyelade, B. I. O. Ade-Omowaye, V. F. Adeomi, Influence of variety on protein, fat contents and some physical characteristics of okra seeds, *Journal of Food Engineering*, 2003, 57 (2): 111-114.
- [35] F. W. Martin, L. Telek, R. Ruberte, A. G. Santiago, Protein, Oil and Gossypol Contents of a Vegetable Curd Made from Okra Seeds, *Journal of Food Science*, 1979, 44 (5): 1517-1519.
- [36] P. U. Rao, Chemical composition and biological evaluation of Okra (*Hibiscus esculentus*) seeds and their kernels, *Plant Foods for Human Nutrition (Formerly Qualitas Plantarum)*, 1985, 35 (4): 389-396.
- [37] M. L. Woolfe, M. F. Chaplin and G. Otchere, Studies on the mucilages extracted from okra fruits (*Hibiscus esculentus* L.) and baobab leaves (*Adansonia digitata* L.), *Journal of the Science of Food and Agriculture*, 1977, 28(6): 519-529.
- [38] F. I. El-Nahry, M. I. El-Ghorab and R. Younes, Nutritive value of local varieties of fresh and sundried okra (*hibiscus esculentus*) pods and seeds, *Plant Foods for Human Nutrition (Formerly Qualitas Plantarum)*, 1978, 28 (3): 227-231.
- [39] P. A. Savello, F. W. Martin and J. M. Hill, Nutritional composition of okra seed meal, *J Agric Food Chem*, 1980, 28 (6): 1163-1166.

Keywords: Diabetes mellitus, endocrine, anti-diabetic, *Abelmoschus esculentus*. 1. Introduction Diabetes mellitus, a leading non-communicable disease with multiple etiologies, affects more than 100 million people worldwide and is considered as one of the five leading causes of death in the world (Zimmet PZ. 1999) [1]. It is a metabolic disorder affecting carbohydrate, fat and protein metabolism. Diabetes Mellitus Diabetes mellitus is a metabolic disease usually characterized by the classic triad of polydipsia, polyuria and polyphagia, consequences of homeostasis disruption due to impaired glucose metabolism. (Bascones-Martinez A et al 2011) [11]. Types of Diabetes mellitus • Type 1 diabetes • Type 2 diabetes • Gestational diabetes. ~ 23 ~. Journal of Medicinal Plants Studies. *Abelmoschus esculentus* (AE), a commonly consumed vegetable, is well-known for its anti-hyperglycemic effects. However, few scientific reports have identified its targets because mucilage increases the difficulty of manipulation. We recently reported extraction steps to obtain subfractions of AE, which were found to attenuate the adverse effects of high glucose and fatty acid in vitro. In this study, we used modified extraction steps and type 2 diabetic rats to explore whether AE subfractions can improve the metabolic disturbances caused by insulin resistance in vivo. Discover a faster, simpler path to publishing in a high-quality journal. PLOS ONE promises fair, rigorous peer review, broad scope, and wide readership – a perfect fit for your research every time.